



Posterior and Posterior Inferior Capsular Shift Protocol

Dr. Michael Chiu, MD
Illinois Bone and Joint Institute
Phone: (847)870-6100
Fax: (847)870-8159

Protection Phase (0-6 weeks):

Precautions

- Postoperative brace in 30-45° abduction, 15° external rotation for 6 weeks
- Brace to be worn at all times (even when sleeping) with the exception of exercise activity and bathing
- No overhead activity
- No flexion for first 6 weeks

Goals:

- Allow/promote healing of repaired posterior capsule
- Initiate early protected ROM
- Retard muscular atrophy
- Decrease pain and inflammation

Weeks 0-4

Exercises

- Gripping exercises with putty
- Active elbow flexion-extension and pronation-supination
- Active ROM cervical spine
- Passive ROM progressing to active-assisted ROM of GH joint
 - o External rotation to 25-30° at 30-45° of abduction
 - o Internal rotation to 15-25° at 30-45° of abduction (begin week three)
- Submaximal pain free shoulder isometrics in the plane of the scapula
 - o Flexion
 - o Abduction
 - o Extension
 - o External rotation
- o **Avoid IR at this point**

Note: In general all exercises begin with one set of 10 repetitions and should increase by one set of 10 repetitions daily as tolerated to five sets of 10 repetitions.

Cryotherapy: Ice after exercises for 20 minutes. Ice up to 20 minutes per hour to control pain and swelling.

Weeks 4-6

Goals

- Gradual increase in ROM
- Normalize arthrokinematics
- Improve strength



- Decrease pain and inflammation

Range of motion exercises

- Active-assisted exercises of GH joint
- External rotation in multiple planes of shoulder abduction (up to 90°)
- Shoulder flexion to tolerance
- Elevation in the plane of the scapula to tolerance
- Shoulder abduction (pure) to 90°
- Internal rotation 35° at 45° of abduction
- Pulleys (AAROM)
- Shoulder elevation in the plane of the scapula to tolerance
- Shoulder flexion to tolerance
- **Gentle** self-capsular stretches as needed/indicated

Gentle Joint Mobilization (Grades I-II) to Re-establish Normal Arthrokinematics

- Scapulothoracic joint
- GH joint (avoid posterior glides)
- SC joint
- AC joint

AROM Exercises

- Active abduction to 90°
- Active external rotation to 90°
- IR to 35°

Strengthening Exercises

- Elbow/wrist progressive resistive exercise program

Conditioning Program For:

- Trunk
- Lower extremities
- Cardiovascular endurance

Decrease Pain and Inflammation

- Ice and modalities prn

Brace

- Discontinue 4-6 weeks post-surgery per physicians instruction

Phase 2: Intermediate Phase (Weeks 6-12)

Goals:

- Full, nonpainful ROM at week eight (patient will not have full IR at this time)
- Normalize arthrokinematics
- Enhance strength
- Improve neuromuscular control

Weeks 6-9

Range of Motion Exercises

- A/AROM to AROM as appropriate
- External rotation to tolerance
- Shoulder abduction to tolerance
- Shoulder flexion to tolerance



- Pulleys: flexion, abduction, and elevation in the plane of the scapula to tolerance

- Internal rotation to no more than 40°

Joint Mobilization

- Continue as above as indicated

Strengthening Exercises

- Initiate IR isometrics in slight ER (do not perform past neutral)

- Initiate Theraband for internal and external rotation at 0° abduction (IR later in the phase)

- Initiate isotonic dumbbell program

- Shoulder abduction

- Shoulder flexion

- Latissimus dorsi

- Rhomboids

- Biceps curl

- Triceps kick-out over table

- Push-ups into wall (serratus anterior)

Weeks 10-12

- Continue all exercises listed above

Initiate

- Active internal rotation at 90° GH abduction with elbow at 90° flexion

- Dumbbell supraspinatus

- Theraband exercises for rhomboids, latissimus dorsi, biceps, and triceps

- Progressive push-ups

Phase 3: Dynamic Strengthening Program (Weeks 12-18)

Criteria for Progression to Phase 3

- Full, nonpainful ROM

- No complaints of pain/tenderness

- Strength 70% of contralateral side

Weeks 13-15

Goals

- Enhance strength, power, and endurance

- Enhance neuromuscular control

Emphasis of Phase 3

- High-speed/high-energy strengthening exercises

- Eccentric training

- Diagonal patterns

Exercises

- Continue internal and external rotation Theraband exercises at 0° abduction (arm at side)

- Theraband for rhomboids

- Theraband for latissimus dorsi

- Theraband for a biceps and triceps

- Continue dumbbell exercises for supraspinatus and deltoid



- Progressive serratus anterior push-up-anterior flexion
- Continue trunk and lower extremity strengthening and conditioning exercises
- Continue self-capsular stretches

Progress to:

- Isotonic shoulder strengthening exercises isolating the rotator cuff-including sidelying external rotation, prone arm raises at 0, 90 & 120°, prone external rotation, and internal rotation at 0 & 90°; progress to standing strengthening exercise once able to tolerate resistance against gravity without substitution
- Progress scapulothoracic/upper back musculature strengthening exercises
- Dynamic stabilization exercises
- Proprioceptive Neuromuscular Facilitation (PNF) exercises

Phase 4: Return to Activity Phase (Weeks 21-28)

Criteria for Progression to Phase 4

- Full ROM
- No pain or tenderness
- Satisfactory clinical examination

Goal

- Progressively increase activities to prepare patient for unrestricted functional return

Exercises

- Continue Theraband, and dumbbell exercises outlined in phase 3
- Continue ROM exercises
- Initiate interval programs between weeks 28 and 32 (if patient is a recreational athlete)
- Continue strengthening exercises for scapular and rotator cuff muscles
- Progress to functional activities needed for ADL's and sport
- Thrower's ten program (see protocol)